

Original citation:

Shi, Q., Zudov, M. A., Morrison, Christopher and Myronov, Maksym. (2015) Transport anisotropy in Ge quantum wells in the absence of quantum oscillations. *Physical Review B (Condensed Matter and Materials Physics)*, 92 (16). 161405(R).

Permanent WRAP URL:

<http://wrap.warwick.ac.uk/78527>

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions. Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Publisher statement:

© 2015 American Physical Society

A note on versions:

The version presented here may differ from the published version or, version of record, if you wish to cite this item you are advised to consult the publisher's version. Please see the 'permanent WRAP URL' above for details on accessing the published version and note that access may require a subscription.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk

Transport anisotropy in Ge quantum wells in the absence of quantum oscillations

Q. Shi,¹ M. A. Zudov,^{1,*} C. Morrison,² and M. Myronov²

¹*School of Physics and Astronomy, University of Minnesota, Minneapolis, Minnesota 55455, USA*

²*Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom*

(Received 8 June 2015; published 14 October 2015)

A recent study of a high-mobility two-dimensional hole gas in a strained Ge quantum well revealed strong transport anisotropy in the quantum Hall regime when the magnetic field was tilted away from the sample normal [Q. Shi, M. A. Zudov, C. Morrison, and M. Myronov, *Phys. Rev. B* **91**, 201301(R) (2015)]. In the present study we demonstrate that the anisotropy persists to such high temperatures and filling factors that quantum oscillations are no longer observed. This finding rules out the formation of a stripe phase as a possible origin for the observed anisotropy. However, we also show that the observed anisotropy is not consistent with other known anisotropies, such as those arising from finite-thickness effects or surface roughness.

DOI: [10.1103/PhysRevB.92.161405](https://doi.org/10.1103/PhysRevB.92.161405)

PACS number(s): 73.43.Qt, 73.63.Hs, 73.40.—c

It is well established that transport properties of two-dimensional (2D) systems could be modified by a pure in-plane magnetic field $B = B_{\parallel}$ for several reasons. First, B_{\parallel} can align the spin of the charge carriers, leading to an increase of the resistivity due to suppression of screening by charged impurities [1,2]. Second, due to a finite thickness of a 2D system, B_{\parallel} distorts the Fermi contour and modifies the scattering rates, also producing positive magnetoresistance [3–6]. Finally, an increase in resistivity with B_{\parallel} could also occur because of interface roughness [7–9], due to local, anisotropic perpendicular magnetic fields. Both finite-thickness and roughness mechanisms imply some anisotropy in the resistivity tensor, albeit with different orientations of the anisotropy axis with respect to B_{\parallel} . The spin-polarization scenario, on the other hand, does not lead to anisotropy, unless the crystal structure is anisotropic. An addition of a weak perpendicular magnetic field ($B_{\perp} \ll B_{\parallel}$) can further modify the in-plane magnetoresistance and anisotropy. However, if one disregards the appearance of quantum oscillations, the effect of B_{\perp} is usually rather small [3,10–14].

In a purely perpendicular magnetic field, $B = B_{\perp}$, 2D systems reveal a much wider variety of transport phenomena. At low B_{\perp} , these phenomena include several kinds of both positive and negative [15–22] magnetoresistances, which can originate from electron-electron interactions [23–26] or quasiclassical memory effects [27–33]. At higher B_{\perp} , much more dramatic phenomena, such as integer [34] and fractional [35] quantum Hall (QH) effects, stripe and bubble phases [36–39], as well as Wigner crystals [40–44], emerge due to the interplay among Landau/Zeman quantizations, disorder, and electron-electron interactions. Added B_{\parallel} can significantly change the transport properties owing to, e.g., spin polarization [45–47], modification of scattering rates [48–50], and finite-thickness effects [51]. Unless already anisotropic, the system remains isotropic with few exceptions, such as a B_{\parallel} -induced stripe phase in the $N = 1$ Landau level [52,53].

It was recently realized that when a high-mobility 2D hole gas (2DHG) in a strained Ge quantum well is subject to *both* in-plane ($B_x = B \sin \theta$) and *out-of-plane* ($B_z = B \cos \theta$) magnetic fields, its low temperature transport properties in the

QH regime become strongly anisotropic [54]. At $T \approx 0.3$ K, and B_z larger than the onset of spin splitting, the resistivity ratio at half-integer filling factors was found to increase gradually with θ , reaching $\rho_{xx}/\rho_{yy} \approx 11.5$ at $\theta = 80^\circ$. At smaller B_z , the anisotropy decreased roughly linearly with B_z for *all* θ , until vanishing close to the onset of Shubnikov–de Haas oscillations. Finally, switching off *either* B_z or B_x resulted in a roughly isotropic state with $\rho_{xx}/\rho_{yy} \approx 1$ over a wide range of B_x or B_z (up to 7 T).

The observed anisotropy was examined in terms of a stripe/nematic phase [36–39], known to occur in high ($2 \leq N \leq 6$) Landau levels of ultraclean GaAs systems cooled down to $T \lesssim 0.1$ K [37,38]. While some features were consistent with the stripe scenario, the slow decay of anisotropy with T seemed to rule against it. As the focus of Ref. [54] was on near half-integer filling factors for $4 < \nu < 40$ in the QH regime, measurements were limited to $T < 1.5$ K and moderate B_x , which implied $B_x/B_z < 6$. It is thus important to investigate if anisotropy can survive at higher T and higher tilt angles when the quantum oscillations are absent. It is also interesting to extend the study to the lower $N < 2$ Landau levels, where the nematic phases in GaAs are less likely to occur.

In this Rapid Communication we report on transport measurements in a high-mobility 2DHG in a Ge quantum well in tilted magnetic fields up to 18 T, focusing on the regime of (i) much higher B_x/B_z and T up to 8 K and (ii) the $N = 1$ Landau level. We find that while the anisotropy smoothly increases with B_x , the addition of a small perpendicular magnetic field $B_z \lesssim 0.5$ T significantly enhances the anisotropy without bringing in quantum oscillations. At $B_z \gtrsim 0.5$ T, we find that the main result of Ref. [54], namely, that ρ_{xx}/ρ_{yy} is determined by the tilt angle alone, holds all the way up to $B_x/B_z \gtrsim 20$ and to much higher T , even in the absence of quantum oscillations. The existence of anisotropy in the regime where no quantum oscillations are seen allows us to rule out the formation of a stripe phase as a possible origin. We further demonstrate that our findings are not compatible with other known anisotropies, such as those arising from finite-thickness effects [2] or surface roughness [7–9], pointing towards a different mechanism of anisotropic transport. We also find that at low temperatures and at fixed tilt angle, anisotropy is significantly suppressed in the $N = 1$ Landau level, indicating that the “scaling” of anisotropy

*Corresponding author: zudov@physics.umn.edu




FIG. 1. (Color online) (a) $R_{xx}(B)$ (in k Ω) and (b) $R_{yy}(B)$ (in Ω) at $\theta = 80^\circ$ (solid lines) and 90° (dashed lines) at $T \approx 0.3$ K. (c) Same as above on a log-linear scale. (d) $\rho_{xx}(B)$, $\rho_{yy}(B)$ and (e) $\rho_{xx}/\rho_{yy}(B)$ at $\theta = 90^\circ$.

with a tilt angle breaks down (see the Supplemental Material [55]).

Our sample is a 5×5 mm square fabricated from a fully strained, 17-nm-wide Ge quantum well grown by reduced pressure chemical vapor deposition on a relaxed $\text{Si}_{0.16}\text{Ge}_{0.84}/\text{Ge}/\text{Si}$ (001) virtual substrate [56–62]. Holes are supplied by a 12-nm-wide B-doped layer separated from the interface by a 30-nm-wide undoped $\text{Si}_{0.16}\text{Ge}_{0.84}$ spacer. At $T = 0.3$ K, our 2DHG has a density $p \approx 2.9 \times 10^{11} \text{ cm}^{-2}$ and mobility $\mu \approx 1.3 \times 10^6 \text{ cm}^2/\text{Vs}$. The resistances R_{xx} and R_{yy} were measured using corner contacts by a low-frequency (a few Hz) lock-in technique. The sample was mounted on a rotator stage and the angle between the sample normal and the magnetic field could be changed *in situ* without warming up the sample. Magnetotransport measurements were done by either sweeping a magnetic field at a fixed angle or rotating the sample in a fixed magnetic field.

In Fig. 1 we compare magnetoresistances $R_{xx}(B)$ [Fig. 1(a)] and $R_{yy}(B)$ [Fig. 1(b)] measured in a parallel field ($\theta = 90^\circ$, $B = B_{\parallel} = B_x$, dashed line) to their values in a tilted field ($\theta = 80^\circ$, $B \approx 1.015 B_x$, solid line). All four traces shown in Figs. 1(a) and 1(b) are also presented in Fig. 1(c) on a log-linear scale. At $B = 0$, our 2DHG exhibits modest anisotropy with $R_{xx} < R_{yy}$, which likely originates from anisotropic surface roughness [63]. This anisotropy virtually disappears upon application of a purely perpendicular magnetic field, $B = B_z > 0.1$ T [54]. If a purely parallel field is applied, $B = B_x$ ($\theta = 90^\circ$), R_{xx} increases, R_{yy} decreases, and at $B_x \approx 6$ T one finds $R_{xx} \approx R_{yy}$. On the other hand, when a small perpendicular field is added ($\theta = 80^\circ$), both R_{xx} and R_{yy} show much bigger changes, starting from $B_x \approx 0.5$ T and differing by three orders of magnitude at $B_x \approx 2.8$ T. This value of B_x corresponds to $B_z = 0.5$ T, marked by a dotted vertical line.




FIG. 2. (Color online) (a) R_{xx} (in k Ω) and (b) R_{yy} (in Ω) measured at $T = 4$ K and different B , as marked, vs B_z , introduced via rotation of the sample.

Since R_{xx} and R_{yy} are measured in a square sample, the decrease of R_{yy} does not necessarily mean a decrease of resistivity ρ_{yy} . Following the results of Ref. [64], we convert R_{xx} , R_{yy} to ρ_{xx} , ρ_{yy} and present the results versus B_x at $\theta = 90^\circ$ in Fig. 1(d). We find that ρ_{yy} increases slower than ρ_{xx} , and the resistivity ratio becomes $\rho_{xx}/\rho_{yy} \approx 1.6$ at $B = B_x = 10$ T, as illustrated in Fig. 1(e). We thus confirm that a purely parallel magnetic field produces only a modest transport anisotropy.

To examine the anisotropy in a regime when quantum oscillations are absent, we perform the transport measurements at an elevated temperature of $T = 4$ K and at large tilt angles. To access the high angle limit, we apply a fixed magnetic field B along the \hat{x} direction and then rotate the sample about the \hat{y} axis to introduce a small perpendicular field B_z . In Fig. 2 we present R_{xx} [Fig. 2(a)] and R_{yy} [Fig. 2(b)], measured at $T = 4$ K, versus B_z , introduced via rotation of the sample in different $B \approx B_x$ from 5 to 18 T, as marked. We observe that R_{xx} (R_{yy}) initially increases (decreases) with B_z and then shows a maximum (minimum) at all B studied. With increasing $B \approx B_x$, the maximum (minimum) becomes higher (lower) and gradually shifts to higher B_z . Based on these observations we conclude that the anisotropy (a) can be significant even at $T = 4$ K, (b) does not require quantum oscillations, and (c) monotonically increases with B_x while exhibiting a maximum at B_z somewhere between 0.2 and 0.8 T.

Having determined the range of B_z where the anisotropy is maximized, we present in Fig. 3 a false color plot of ρ_{xx}/ρ_{yy}




FIG. 3. (Color online) ρ_{xx}/ρ_{yy} vs (B_z, B_x) at $T = 4$ K. Dotted lines are drawn at $B_x/B_z = 5, 10, 15$, and 30 .

vs B_z and B_x . The strongest anisotropy, characterized by $\rho_{xx}/\rho_{yy} > 10$, occurs at $B_x \gtrsim 10$ T in a region which is domed at $B_z \approx 0.5$ T. This dome has a considerably larger gradient on the lower B_z side than at the higher B_z side. Furthermore, the iso-anisotropy lines on the higher B_z side are well described by constant B_x/B_z , as illustrated by dotted lines. While this result was already obtained for half-integer filling factors in the QH regime at $T \approx 0.3$ K and $B_x/B_z < 5.7$ [54], here we demonstrate that the same rule applies for much higher T , in the regime where there are no quantum oscillations, and up to much higher B_x/B_z . However, this rule breaks down on the other side of the dome, where, as we show next, the anisotropy is controlled by the perpendicular component of the magnetic field.

In Fig. 4(a) we present ρ_{xx}/ρ_{yy} vs B_z measured at different $B_x \approx B$, as marked. At small B_z , ρ_{xx}/ρ_{yy} shows a roughly linear increase with approximately the same slope for all B , which culminates with a maximum at $B_z \approx 0.5$ T. In Fig. 4(b) we replot the same data versus B_z/B and observe that the




FIG. 4. (Color online) ρ_{xx}/ρ_{yy} vs (a) B_z and (b) B_z/B at $B_x \approx B = 5, 8, 11, 14$, and 18 T at $T = 4$ K.




FIG. 5. (Color online) (a) R_{xx} (in kΩ) and (b) R_{yy} (in Ω) at $B_x \approx B = 8$ T vs B_z , introduced via rotation of the sample, at $T = 0.3, 4$, and 8 K. (c) ρ_{xx}/ρ_{yy} vs T at $\theta = 60^\circ, 72^\circ, 86^\circ$, and 90° , as marked. Dotted lines are guides for the eye.

decreasing parts of all curves collapse onto one. Consistent with Ref. [54], studying half-integer filling factors in the QH regime, the observed collapse once again confirms that in this parameter range the anisotropy is determined only by the tilt angle.

To examine the temperature dependence of the anisotropy in this regime, in Fig. 5 we present R_{xx} [Fig. 5(a)] and R_{yy} [Fig. 5(b)] vs B_z measured at $B_x \approx B = 8$ T and $T = 0.3, 4$, and 8 K. With increasing temperature, R_{xx} decreases while R_{yy} increases, signaling a decrease in anisotropy over the whole range of B_z , except $B_z = 0$. In Fig. 4(c) we present ρ_{xx}/ρ_{yy} vs T , measured at $\theta = 60^\circ, 72^\circ, 86^\circ$, and 90° , as marked. We observe that at all tilt angles (except $\theta = 90^\circ$), the anisotropy decays with increasing temperature and that the rate of this decay drops considerably with increasing tilt angle. Indeed, while at $\theta = 60^\circ$, the anisotropy disappears at $T \approx 2$ K, the resistivity ratio measured at $\theta = 86^\circ$ remains significant, and $\rho_{xx}/\rho_{yy} > 4$, even at $T = 8$ K. At $\theta = 90^\circ$, on the other hand, we observe virtually no temperature dependence of the anisotropy. This finding suggests that the mechanism responsible for the temperature dependence in tilted fields is completely absent in pure $B_{||}$.

The observation of strong anisotropy in small B_z and at high temperatures unambiguously rules out QH stripes as a possible origin. First, the robustness against temperature suggests a much larger energy scale than expected of the charge density wave. In the Hartree-Fock approach, the latter scales with the exchange energy [36,65], and is ~ 1 K at $B_z = 0.5$ T. Indeed, in clean GaAs systems, stripes manifest only at much lower T , even in tilted magnetic fields. Second, QH stripes are expected only when spin splitting is resolved, while in our experiment, at large enough $B_{||}$, the anisotropy sets in as soon as B_z is added.

The perpendicular magnetic field certainly plays a crucial role in the underlying mechanism of the anisotropy. Our Ge quantum wells exhibit a modest transport anisotropy both at $B = 0$ and in a purely in-plane magnetic field [see Fig. 1(e) and Ref. [54]]. It is known that an in-plane field could induce anisotropy due to the distortion of the Fermi contour [3,4] and surface roughness, via anisotropic, random perpendicular magnetic fields [7–9]. While the former can be ruled out because it leads to $\rho_{xx} < \rho_{yy}$ (when $B_{\parallel} = B_x$), the latter is consistent with our observations. We thus conclude that these modest anisotropies likely originate from the surface roughness [63].

One important question is whether the surface roughness can also result in huge anisotropy in our Ge quantum wells in tilted B . First, B_{\parallel} -induced anisotropy is known to be temperature independent [7,9], whereas the anisotropy in tilted fields has a significant temperature dependence (see Fig. 5 and Ref. [54]). Second, because of the anisotropy at $B = 0$, the magnitude of B_{\parallel} -induced anisotropy must depend on the orientation of B_{\parallel} [9], whereas the observed anisotropy in tilted magnetic fields was found to be insensitive to the orientation of B_{\parallel} [54]. Finally, no strong enhancement of anisotropy due to additional B_z has been reported in experiments using GaAs samples with a much larger surface roughness [7] or predicted theoretically [66–68]. In fact, Ref. [7] reported a *reduction* of the anisotropy upon introduction of B_z . We therefore conclude that the anisotropy in a purely parallel magnetic field is unlikely to be related to the strong anisotropy in tilted magnetic fields.

In summary, we have investigated anisotropic transport in a high-mobility 2D hole gas in a strained Ge quantum well

in tilted magnetic fields up to 18 T and at temperatures up to 8 K. We have found that the maximum of ρ_{xx}/ρ_{yy} occurs at the highest available B_x and at $B_z \approx 0.5$ T, where it remains significant even at the highest temperature studied. The existence of anisotropy in a regime where no quantum oscillations are seen rules out the formation of stripes as a possible origin. Even though quantum oscillations are not required, the perpendicular magnetic field plays a crucial role both in the magnitude of the anisotropy and its temperature dependence. We have also shown that our findings are not compatible with other known anisotropies, such as those arising from finite-thickness effects or surface roughness, suggesting a different mechanism of anisotropic transport.

We thank G. Jones, S. Hannas, T. Murphy, J. Park, and D. Smirnov for technical assistance, and A. MacDonald, L. Engel, S. Kivelson, D. Polyakov, and B. Shklovskii for discussions. The work at Minnesota was funded by the U.S. Department of Energy, Office of Basic Energy Sciences, under Grant No. ER 46640 SC0002567. The work at Warwick was supported in part by EPSRC-funded Spintronic device physics in Si/Ge Heterostructures EP/J003263/1 and Platform Grant No. EP/J001074/1 projects. Q.S. acknowledges support from an Allen M. Goldman fellowship and a University of Minnesota Doctoral Dissertation Fellowship. Experiments were performed at the National High Magnetic Field Laboratory, which is supported by NSF Cooperative Agreement No. DMR-0654118, by the State of Florida, and by the DOE.

-
- [1] E. Tutuc, S. Melinte, and M. Shayegan, *Phys. Rev. Lett.* **88**, 036805 (2002).
 - [2] S. Das Sarma and E. H. Hwang, *Phys. Rev. B* **72**, 035311 (2005).
 - [3] J. M. Heisz and E. Zaremba, *Phys. Rev. B* **53**, 13594 (1996).
 - [4] S. Das Sarma and E. H. Hwang, *Phys. Rev. Lett.* **84**, 5596 (2000).
 - [5] S. J. Papadakis, E. P. De Poortere, M. Shayegan, and R. Winkler, *Phys. Rev. Lett.* **84**, 5592 (2000).
 - [6] X. Zhou, B. A. Piot, M. Bonin, L. W. Engel, S. Das Sarma, G. Gervais, L. N. Pfeiffer, and K. W. West, *Phys. Rev. Lett.* **104**, 216801 (2010).
 - [7] A. A. Bykov, G. M. Gusev, J. R. Leite, A. K. Bakarov, A. V. Goran, V. M. Kudryashev, and A. I. Toropov, *Phys. Rev. B* **65**, 035302 (2001).
 - [8] A. Bykov, A. Bakarov, A. Goran, A. Latyshev, and A. Toropov, *JETP Lett.* **74**, 164 (2001).
 - [9] A. V. Goran, A. A. Bykov, and A. I. Toropov, *Semicond. Sci. Technol.* **23**, 105017 (2008).
 - [10] T. Englert, J. Maan, D. Tsui, and A. Gossard, *Solid State Commun.* **45**, 989 (1983).
 - [11] P. Štředa, P. Vašek, and M. Cukr, *Phys. Rev. B* **51**, 11144 (1995).
 - [12] D. Kamburov, M. Shayegan, R. Winkler, L. N. Pfeiffer, K. W. West, and K. W. Baldwin, *Phys. Rev. B* **86**, 241302 (2012).
 - [13] A. Y. Kuntsevich, L. A. Morgun, and V. M. Pudalov, *Phys. Rev. B* **87**, 205406 (2013).
 - [14] D. Kamburov, M. A. Mueed, M. Shayegan, L. N. Pfeiffer, K. W. West, K. W. Baldwin, J. J. D. Lee, and R. Winkler, *Phys. Rev. B* **88**, 125435 (2013).
 - [15] M. A. Paalanen, D. C. Tsui, and J. C. M. Hwang, *Phys. Rev. Lett.* **51**, 2226 (1983).
 - [16] K. K. Choi, D. C. Tsui, and S. C. Palmateer, *Phys. Rev. B* **33**, 8216 (1986).
 - [17] L. Li, Y. Y. Proskuryakov, A. K. Savchenko, E. H. Linfield, and D. A. Ritchie, *Phys. Rev. Lett.* **90**, 076802 (2003).
 - [18] L. Bockhorn, P. Barthold, D. Schuh, W. Wegscheider, and R. J. Haug, *Phys. Rev. B* **83**, 113301 (2011).
 - [19] A. T. Hatke, M. A. Zudov, J. L. Reno, L. N. Pfeiffer, and K. W. West, *Phys. Rev. B* **85**, 081304(R) (2012).
 - [20] Q. Shi, P. D. Martin, Q. A. Ebner, M. A. Zudov, L. N. Pfeiffer, and K. W. West, *Phys. Rev. B* **89**, 201301(R) (2014).
 - [21] Q. Shi, M. A. Zudov, L. N. Pfeiffer, and K. W. West, *Phys. Rev. B* **90**, 201301(R) (2014).
 - [22] L. Bockhorn, I. V. Gornyi, D. Schuh, C. Reichl, W. Wegscheider, and R. J. Haug, *Phys. Rev. B* **90**, 165434 (2014).
 - [23] B. L. Altshuler and A. G. Aronov, in *Electron-Electron Interactions in Disordered Systems*, edited by A. L. Efros and M. Pollak (North-Holland, Amsterdam, 1985).
 - [24] S. M. Girvin, M. Jonson, and P. A. Lee, *Phys. Rev. B* **26**, 1651 (1982).
 - [25] I. V. Gornyi and A. D. Mirlin, *Phys. Rev. Lett.* **90**, 076801 (2003).

- [26] I. V. Gornyi and A. D. Mirlin, *Phys. Rev. B* **69**, 045313 (2004).
- [27] E. M. Baskin, L. N. Magarill, and M. V. Entin, *Sov. Phys. JETP* **48**, 365 (1978).
- [28] A. V. Bobylev, F. A. Maaø, A. Hansen, and E. H. Hauge, *Phys. Rev. Lett.* **75**, 197 (1995).
- [29] M. M. Fogler, A. Y. Dobin, V. I. Perel, and B. I. Shklovskii, *Phys. Rev. B* **56**, 6823 (1997).
- [30] A. Dmitriev, M. Dyakonov, and R. Jullien, *Phys. Rev. B* **64**, 233321 (2001).
- [31] A. Dmitriev, M. Dyakonov, and R. Jullien, *Phys. Rev. Lett.* **89**, 266804 (2002).
- [32] A. D. Mirlin, D. G. Polyakov, F. Evers, and P. Wölffe, *Phys. Rev. Lett.* **87**, 126805 (2001).
- [33] D. G. Polyakov, F. Evers, A. D. Mirlin, and P. Wölffe, *Phys. Rev. B* **64**, 205306 (2001).
- [34] K. von Klitzing, G. Dorda, and M. Pepper, *Phys. Rev. Lett.* **45**, 494 (1980).
- [35] D. C. Tsui, H. L. Stormer, and A. C. Gossard, *Phys. Rev. B* **25**, 1405 (1982).
- [36] A. A. Koulakov, M. M. Fogler, and B. I. Shklovskii, *Phys. Rev. Lett.* **76**, 499 (1996).
- [37] M. P. Lilly, K. B. Cooper, J. P. Eisenstein, L. N. Pfeiffer, and K. W. West, *Phys. Rev. Lett.* **82**, 394 (1999).
- [38] R. R. Du, D. C. Tsui, H. L. Stormer, L. N. Pfeiffer, K. W. Baldwin, and K. W. West, *Solid State Commun.* **109**, 389 (1999).
- [39] E. Fradkin and S. A. Kivelson, *Phys. Rev. B* **59**, 8065 (1999).
- [40] E. Wigner, *Phys. Rev.* **46**, 1002 (1934).
- [41] Y. E. Lozovik and V. I. Yudson, *JETP Lett.* **22**, 11 (1975).
- [42] P. K. Lam and S. M. Girvin, *Phys. Rev. B* **30**, 473 (1984).
- [43] D. Levesque, J. J. Weis, and A. H. MacDonald, *Phys. Rev. B* **30**, 1056 (1984).
- [44] Y. P. Chen, G. Sambandamurthy, Z. H. Wang, R. M. Lewis, L. W. Engel, D. C. Tsui, P. D. Ye, L. N. Pfeiffer, and K. W. West, *Nat. Phys.* **2**, 452 (2006).
- [45] R. J. Nicholas, R. J. Haug, K. v. Klitzing, and G. Weimann, *Phys. Rev. B* **37**, 1294 (1988).
- [46] R. R. Du, H. L. Stormer, D. C. Tsui, L. N. Pfeiffer, and K. W. West, *Phys. Rev. Lett.* **70**, 2944 (1993).
- [47] D. R. Leadley, R. J. Nicholas, J. J. Harris, and C. T. Foxon, *Phys. Rev. B* **58**, 13036 (1998).
- [48] G. M. Gusev, J. R. Leite, A. A. Bykov, N. T. Moshegov, V. M. Kudryashev, A. I. Toropov, and Y. V. Nastaushchev, *Phys. Rev. B* **59**, 5711 (1999).
- [49] A. T. Hatke, M. A. Zudov, L. N. Pfeiffer, and K. W. West, *Phys. Rev. B* **83**, 081301(R) (2011).
- [50] A. Bogan, A. T. Hatke, S. A. Studenikin, A. Sachrajda, M. A. Zudov, L. N. Pfeiffer, and K. W. West, *Phys. Rev. B* **86**, 235305 (2012).
- [51] A. T. Hatke, M. A. Zudov, L. N. Pfeiffer, and K. W. West, *Phys. Rev. B* **85**, 241305(R) (2012).
- [52] M. P. Lilly, K. B. Cooper, J. P. Eisenstein, L. N. Pfeiffer, and K. W. West, *Phys. Rev. Lett.* **83**, 824 (1999).
- [53] W. Pan, R. R. Du, H. L. Stormer, D. C. Tsui, L. N. Pfeiffer, K. W. Baldwin, and K. W. West, *Phys. Rev. Lett.* **83**, 820 (1999).
- [54] Q. Shi, M. A. Zudov, C. Morrison, and M. Myronov, *Phys. Rev. B* **91**, 201301(R) (2015).
- [55] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevB.92.161405> for transport measurements in the $N = 1$ Landau level.
- [56] A. Dobbie, M. Myronov, R. J. H. Morris, A. H. A. Hassan, M. J. Prest, V. A. Shah, E. H. C. Parker, T. E. Whall, and D. R. Leadley, *Appl. Phys. Lett.* **101**, 172108 (2012).
- [57] M. A. Zudov, O. A. Mironov, Q. A. Ebner, P. D. Martin, Q. Shi, and D. R. Leadley, *Phys. Rev. B* **89**, 125401 (2014).
- [58] C. Morrison, P. Wiśniewski, S. D. Rhead, J. Foronda, D. R. Leadley, and M. Myronov, *Appl. Phys. Lett.* **105**, 182401 (2014).
- [59] M. Myronov, C. Morrison, J. Halpin, S. Rhead, C. Casteleiro, J. Foronda, V. A. Shah, and D. Leadley, *Jpn. J. Appl. Phys.* **53**, 04EH02 (2014).
- [60] Q. Shi, Q. A. Ebner, and M. A. Zudov, *Phys. Rev. B* **90**, 161301(R) (2014).
- [61] M. Myronov, C. Morrison, J. Halpin, S. Rhead, J. Foronda, and D. Leadley, *Solid-State Electron.* **110**, 35 (2015).
- [62] Q. Shi, M. A. Zudov, C. Morrison, and M. Myronov, *Phys. Rev. B* **91**, 241303 (2015).
- [63] A. H. A. Hassan, R. J. H. Morris, O. A. Mironov, R. Beanland, D. Walker, S. Huband, A. Dobbie, M. Myronov, and D. R. Leadley, *Appl. Phys. Lett.* **104**, 132108 (2014).
- [64] S. H. Simon, *Phys. Rev. Lett.* **83**, 4223 (1999).
- [65] M. M. Fogler, A. A. Koulakov, and B. I. Shklovskii, *Phys. Rev. B* **54**, 1853 (1996).
- [66] A. D. Mirlin, D. G. Polyakov, and P. Wölffe, *Phys. Rev. Lett.* **80**, 2429 (1998).
- [67] M. Calvo, *Phys. Rev. B* **57**, R4241 (1998).
- [68] F. Evers, A. D. Mirlin, D. G. Polyakov, and P. Wölffe, *Phys. Rev. B* **60**, 8951 (1999).